
Newton's First & Second Law

AP Physics C

Facts about FORCE

- Unit is the **NEWTON(N)**
 - Is by definition a push or a pull
 - Can exist during physical contact(Tension, Friction, Applied Force)
 - Can exist with NO physical contact, called FIELD FORCES (gravitational, electric, etc)
-

Newton's First Law – The Law of Inertia

INERTIA – a quantity of matter, also called **MASS**. Italian for “LAZY”. Unit for MASS = **kilogram**.

Weight or Force due to Gravity is how your MASS is effected by gravity.

$$W = mg$$

NOTE: MASS and WEIGHT are NOT the same thing. MASS never changes When an object moves to a different planet.

What is the weight of an 85.3-kg person on earth? On Mars=3.2 m/s/s)?

$$W = mg \rightarrow W = (85.3)(9.8) = 835.94N$$

$$W_{MARS} = (85.3)(3.2) = 272.96N$$

Newton's First Law

An object in motion remains in motion in a straight line and at a constant speed **OR** an object at rest remains at rest, **UNLESS** acted upon by an **EXTERNAL** (unbalanced) Force.

There are **TWO** conditions here and one constraint.

Condition #1 - The object **CAN** move but must be at a **CONSTANT SPEED**

Condition #2 - The object is at **REST**


Constraint - As long as the forces are **BALANCED!!!!** And if all the forces are balanced the **SUM** of all the forces is **ZERO**.

The bottom line: There is **NO ACCELERATION** in this case **AND** the object must be at **EQUILIBRIUM** (All the forces cancel out).

$$acc = 0 \rightarrow \sum F = 0$$


Free Body Diagrams

A pictorial representation of forces complete with labels.



- **Weight (mg)** - Always drawn from the center, straight down
- **Force Normal (F_N)** - A surface force always drawn perpendicular to a surface.
- **Tension (T or F_T)** - force in ropes and always drawn AWAY from object.
- **Friction (F_f)** - Always drawn opposing the motion.

Free Body Diagrams



N.F.L and Equilibrium

Since the $F_{\text{net}} = 0$, a system moving at a constant speed or at rest MUST be at **EQUILIBRIUM.**


TIPS for solving problems

- Draw a FBD
 - Resolve anything into COMPONENTS
 - Write equations of equilibrium
 - Solve for unknowns
-

Example

A 10-kg box is being pulled across the table to the right at a constant speed with a force of 50N.

- Calculate the Force of Friction $F_a = F_f = 50N$
- Calculate the Force Normal



$$mg = F_n = (10)(9.8) = 98N$$

Example


Suppose the same box is now pulled at an angle of 30 degrees above the horizontal.

a) Calculate the Force of Friction

$$F_{ax} = F_a \cos \theta = 50 \cos 30 = 43.3N$$

$$F_f = F_{ax} = 43.3N$$

b) Calculate the Force Normal



$$F_N \neq mg!$$

$$F_N + F_{ay} = mg$$

$$F_N = mg - F_{ay} \rightarrow (10)(9.8) - 50 \sin 30$$

$$F_N = 73N$$

What if it is NOT at Equilibrium?

If an object is NOT at rest or moving at a constant speed, that means the FORCES are UNBALANCED. One force(s) in a certain direction over power the others.

THE OBJECT WILL THEN ACCELERATE.

Newton's Second Law

The acceleration of an object is directly proportional to the NET FORCE and inversely proportional to the mass.

$$a \propto F_{NET} \quad a \propto \frac{1}{m}$$

$$a = \frac{F_{NET}}{m} \rightarrow F_{NET} = ma$$


$$F_{NET} = \sum F$$

Tips:

- Draw an FBD
- Resolve vectors into components
- Write equations of motion by adding and subtracting vectors to find the NET FORCE. Always write larger force - smaller force.
- Solve for any unknowns

N.S.L

A 10-kg box is being pulled across the table to the right by a rope with an applied force of 50N. Calculate the acceleration of the box if a 12 N frictional force acts upon it.



In which direction, is this object accelerating?

The X direction!

So N.S.L. is worked out using the forces in the "x" direction only

$$F_{Net} = ma$$


$$F_a - F_f = ma$$

$$50 - 12 = 10a$$

$$a = 3.8 \text{ m/s}^2$$

Example

A mass, $m_1 = 3.00\text{kg}$, is resting on a frictionless horizontal table is connected to a cable that passes over a pulley and then is fastened to a hanging mass, $m_2 = 11.0\text{ kg}$ as shown below. Find the acceleration of each mass and the tension in the cable.



$$F_{Net} = ma$$

$$m_2g - T = m_2a$$

$$T = m_1a$$

$$m_2g - m_1a = m_2a$$

$$m_2g = m_2a + m_1a$$

$$m_2g = a(m_2 + m_1)$$

$$a = \frac{m_2g}{m_1 + m_2} \rightarrow \frac{(11)(9.8)}{14} = 7.7 \text{ m/s}^2$$

Example

$$F_{Net} = ma$$


$$m_2g - T = m_2a$$

$$T = m_1a$$

$$T = (3)(7.7) = 23.1 N$$

$$F_{Net} = ma \rightarrow \frac{F_{NET}}{a} = m$$

$$Slope = \frac{Rise}{Run}$$



Where does the calculus fit in?

$$\vec{F} = m\vec{a} = m \frac{dv}{dt} = m \frac{d^2x}{dt^2}$$

There could be situations where you are given a displacement function or velocity function. The derivative will need to be taken once or twice in order to get the acceleration. Here is an example.

You are standing on a bathroom scale in an elevator in a tall building. Your mass is 72-kg. The elevator starts from rest and travels upward with a speed that varies with time according to:

$$v(t) = 3t + 0.20t^2$$

When $t = 4.0\text{s}$, what is the reading on the bathroom scale (a.k.a. Force Normal)?

$$a = \frac{dv}{dt} = \frac{d(3t + 0.20t^2)}{dt} = 3 + 0.40t$$

$$a(4) = 3 + 0.40(4) = \mathbf{4.6 \text{ m/s/s}}$$

$$F_{net} = ma$$

$$F_N - mg = ma \rightarrow F_N = ma + mg$$

$$F_N = (72)(9.8) + (72)(4.6) =$$

$$\mathbf{1036.8 \text{ N}}$$

